


SCAI EXPERT CONSENSUS ALGORITHM ON THE MANAGEMENT OF CALCIFIED CORONARY LESIONS¹




SCAI RECOMMENDATIONS FOR IVL IN CORONARY CALCIUM



Consider IVL for concentric, eccentric, and nodular lesions and in settings where atherectomy may be relatively contraindicated



IVL can be used synergistically with atherectomy devices, especially in balloon uncrossable lesions or longer lesions where there is often more heterogeneity in vessel size and pattern of calcification



IVL can be used with multiple guide wires in place (bifurcation lesions)




Intravascular imaging is useful determining an IVL pulse strategy



Intravascular imaging demonstrates that IVL results in both superficial and deep, radial and longitudinal, macrofractures, and microfractures

TRENDS IN USE OF CALCIUM MODIFICATION TOOLS AMONG PATIENTS UNDERGOING PCI²



“

Interestingly, since the approval of IVL in the United States, the frequency of advanced lesion preparation has increased by 40%, while the use of atherectomy has remained relatively stable, suggesting that IVL has facilitated more widespread adoption of advanced lesion preparation³

”

THESE TWO PUBLICATIONS...

- Highlight the growing clinical applications of coronary IVL, with an increasing role as the front-line calcium modification tool for balloon crossable lesions.
- Reinforce the proven safety and efficacy of coronary IVL across all types of calcium morphologies: concentric, eccentric, and nodular.
- Emphasize IVL's unmatched safety profile due to its unique mechanism-of-action that differentiates it from other calcium modification technologies.

1. Riley R.F. Patel M.P. Abbott J.D. et al. SCAI expert consensus statement on the management of calcified coronary lesions. J Soc Cardiovasc Angiogr Interv. 2024.
 2. Butala N. M. Waldo S. W. et al. Use of calcium modification during percutaneous coronary intervention after introduction of coronary intravascular lithotripsy. J Soc Cardiovasc Interv. 2024.
 3. Ali Z. A. Shin D. et al. Between a rock and a hard place: consensus statement on calcified coronary lesions. J Soc Cardiovasc Interv. 2024.

In the United States: Rx only. Indications for Use — The Shockwave Intravascular Lithotripsy (IVL) System with the Shockwave C²⁺ Coronary IVL Catheter is indicated for lithotripsy-enabled, low-pressure balloon dilatation of severely calcified, stenotic de novo coronary arteries prior to stenting. **Contraindications** — The Shockwave C²⁺ Coronary IVL System is contraindicated for the following: This device is not intended for stent delivery. This device is not intended for use in carotid or cerebrovascular arteries. **Warnings** — Use the IVL Generator in accordance with recommended settings as stated in the Operator's Manual. The risk of a dissection or perforation is increased in severely calcified lesions undergoing percutaneous treatment, including IVL. Appropriate provisional interventions should be readily available. Balloon loss of pressure was associated with a numerical increase in dissection which was not statistically significant and was not associated with MACE. Analysis indicates calcium length is a predictor of dissection and balloon loss of pressure. IVL generates mechanical pulses which may cause atrial or ventricular capture in bradycardic patients. In patients with implantable pacemakers and defibrillators, the asynchronous capture may interact with the sensing capabilities. Monitoring of the electrocardiographic rhythm and continuous arterial pressure during IVL treatment is required. In the event of clinically significant hemodynamic effects, temporarily cease delivery of IVL therapy. **Precautions** — Only to be used by physicians trained in angiography and intravascular coronary procedures. Use only the recommended balloon inflation medium. Hydrophilic coating to be wet only with normal saline or water and care must be taken with sharp objects to avoid damage to the hydrophilic coating. Appropriate anticoagulant therapy should be administered by the physician. Precaution should be taken when treating patients with previous stenting within 5mm of target lesion. **Potential adverse effects consistent with standard based cardiac interventions include** — Abrupt vessel closure - Allergic reaction to contrast medium, anticoagulant and/or antithrombotic therapy-Aneurysm-Arrhythmia-Arteriovenous fistula-Bleeding complications -Cardiac tamponade or pericardial effusion-Cardiopulmonary arrest-Cerebrovascular accident (CVA)-Coronary artery/vessel occlusion, perforation, or dissection-Coronary artery spasm-Death-Bleeding complications -Fracture of the guide wire or failure/malfunction of any component of the device that may or may not lead to device embolism, dissection, serious injury or surgical intervention-Hematoma at the vascular access site(s)- Hemorrhage-Hypertension/Hypotension-Infection/sepsis/fever- Myocardial Infarction-Myocardial Ischemia or unstable angina Pain-Peripheral Ischemia-Pseudoaneurysm-Renal failure/insufficiency-Restenosis of the treated coronary artery leading to revascularization-Shock/pulmonary edema-Slow flow, no reflow, or abrupt closure of coronary artery-Stroke Thrombus-Vessel closure, abrupt-Vessel injury requiring surgical repair-Vessel dissection, perforation, rupture, or spasm. **Risks identified as related to the device and its use** — Allergic/immunologic reaction to the catheter material(s) or coating-Device malfunction, failure, or balloon loss of pressure leading to device embolism, dissection, serious injury or surgical intervention-Atrial or ventricular extrasystole-Atrial or ventricular capture. **Prior to use, please reference the Instructions for Use for more information on indications, contraindications, warnings, precautions and adverse events.**
www.shockwavemedical.com/IFU. Please contact your local Shockwave representative for specific country availability and refer to the Shockwave C²⁺ instructions for use containing important safety information.